arXiv:math/9409228v1 [math.CA] 6 Sep 1994

Stieltjes meeting, Delft, 31 Oct.-4 Nov. 1994

Problem

for Mathematicaists, Reducectionists, Macsymalists, and other Mapleators

Painlevé equations for semi-classical recurrence coefficients.

Alphonse P. Magnus
Institut Mathématique, Université Catholique de Louvain, Chemin du Cyclotron 2, B-1348 Louvain-la-Neuve, Belgium.
e-mail: `magnus@anma.ucl.ac.be`

Background: orthonormal polynomials $p_n(x) = \gamma_n x^n + \cdots$:

$$a_{n+1}p_{n+1}(x) = (x-b_n)p_n(x) - a_n p_{n-1}(x) \tag{1}$$

related to a semi-classical weight w : $$\frac{w'(x)}{w(x)} = \frac{2V(x)}{W(x)} \tag{2}$$
where V and W are polynomials, satisfy differential relations

$$Wp_n' = (\Omega_n - V)p_n - a_n\Theta_n p_{n-1} \tag{3}$$

where Ω_n and Θ_n are polynomials of degrees $\leq m-1$ and $m-2$ (if degrees V and $W \leq m-1$ and m), with $$\Omega_{n+1}(x) = (x-b_n)\Theta_n(x) - \Omega_n(x), \tag{4}$$

$$(x-b_n)(\Omega_{n+1}(x) - \Omega_n(x)) = W(x) + a_{n+1}^2\Theta_{n+1}(x) - a_n^2\Theta_n(x)$$

representing nonlinear recurrence relations for the a_n's and b_n's [Laguerre, 19th century].

Statement: *if, in addition to* (2), *the weight* w *depends on a parameter* t *so that*

$$\frac{\partial w(x,t)/\partial t}{w(x,t)} = \frac{T(x,t)}{W(x,t)}, \tag{5}$$

where T *and* W *are polynomials in* x *and* t (*the same* W, *which will now be written* $W(x,t)$), *each recurrence coefficient* $a_n^2(t)$ *and* $b_n(t)$ *satisfy a nonlinear differential equation having Painlevé property* (*movable singular points are only poles*) . [many people, 20th century].

I have an (incomplete) elementary proof of this for **generalized Jacobi weights**

$$\left.\begin{aligned} w(x,t) &= 0, \qquad x < x_1(t) \text{ or } x > x_m(t) \\ &= C_j \prod_{k=1}^{m} |x - x_k(t)|^{\alpha_k}, \quad x_j(t) < x < x_{j+1}(t),\ \ j = 1,\ldots,m-1. \end{aligned}\right\} \tag{6}$$

where therefore $W(x,t) = \prod_1^m (x - x_k(t))$ and (2) and (5) become

$$\frac{w'(x,t)}{w(x,t)} = \sum_{k=1}^{m} \frac{\alpha_k}{x - x_k(t)} \text{ and } \frac{\partial w(x,t)/\partial t}{w(x,t)} = -\sum_{k=1}^{m} \frac{\alpha_k \dot{x}_k(t)}{x - x_k(t)} \tag{7}$$

(N.B. so, the C_j's and the α_j's do *not* depend on t; remark also that $\alpha_k = 2V(x_k)/W'(x_k)$, $\Omega_n(x) = (n+(\sum_1^m \alpha_k)/2)x^{m-1}+\cdots$, $\Theta_n(x) = (2n+1+\sum_1^m \alpha_k)x^{m-2}+\cdots)$. A differential system involving a_n and b_n is then

$$\frac{\dot a_n}{a_n} = \frac{1}{2}\sum_{k=1}^{m} \frac{(\Theta_n(x_k)-\Theta_{n-1}(x_k))\dot x_k}{W'(x_k)}, \tag{8}$$

$$\dot b_n = \sum_{k=1}^{m} \frac{(\Omega_{n+1}(x_k)-\Omega_n(x_k))\dot x_k}{W'(x_k)} = \sum_{k=1}^{m} \frac{((x_k-b_n)\Theta_n(x_k)-2\Omega_n(x_k))\dot x_k}{W'(x_k)}, \tag{9}$$

$$\frac{\dot\gamma_n}{\gamma_n} = -\frac{1}{2}\sum_{k=1}^{m} \dot x_k \Theta_n(x_k)/W'(x_k) \tag{10}$$

$$\frac{d}{dt}\frac{\Theta_n(x_j)}{W'(x_j)} = 2\frac{\dot\gamma_n}{\gamma_n}\frac{\Theta_n(x_j)}{W'(x_j)} - 2\sum_{k\neq j}\frac{\dot x_j-\dot x_k}{x_j-x_k}\frac{\Theta_n(x_k)\Omega_n(x_j)-\Theta_n(x_j)\Omega_n(x_k)}{W'(x_j)W'(x_k)}, \tag{11}$$

$$\frac{d}{dt}\frac{\Theta_{n-1}(x_j)}{W'(x_j)} = -2\frac{\dot\gamma_{n-1}}{\gamma_{n-1}}\frac{\Theta_{n-1}(x_j)}{W'(x_j)} + 2\sum_{k\neq j}\frac{\dot x_j-\dot x_k}{x_j-x_k}\frac{\Theta_{n-1}(x_k)\Omega_n(x_j)-\Theta_{n-1}(x_j)\Omega_n(x_k)}{W'(x_j)W'(x_k)}, \tag{12}$$

$$\frac{d}{dt}\frac{\Omega_n(x_j)}{W'(x_j)} = a_n^2\sum_{k\neq j}\frac{\dot x_j-\dot x_k}{x_j-x_k}\frac{\Theta_n(x_j)\Theta_{n-1}(x_k)-\Theta_n(x_k)\Theta_{n-1}(x_j)}{W'(x_j)W'(x_k)}, \tag{13}$$

Problem: eliminate the Θ's and the Ω's in $(8-13)$ so to exhibit the scalar differential equations for $a_n(t)$ and $b_n(t)$, at least in the $m=3$ case (W of degree 3 in x and 1 in t), and in the even $m=5$ case ($w(x,t) = |x|^\alpha |x^2-x_1^2|^\beta |x^2-x_2^2|^\gamma$, $b_n = 0$), see [1] for confluent cases and [2] for the even case.

Proof of $(8-13)$

The proof is valid only when $\alpha_k > 0$, so that

$$q_n(x) = \int_{\mathbb{R}} \frac{w(u)p_n(u)}{x-u}du \tag{14}$$

is finite at the x_k's. Then, $\frac{d}{dt}\int_{\mathbb{R}} w(u,t)p_n(u,t)p_{n-i}(u,t)du = 0, \quad i = 0,1,\ldots$:

$$\sum_{k=1}^{m}\alpha_k \dot x_k q_n(x_k)p_{n-i}(x_k) + \int_{\mathbb{R}} w(u,t)\frac{\partial p_n(u,t)}{\partial t}p_{n-i}(u,t)du + \frac{\dot\gamma_n}{\gamma_n}\delta_{i,0} = 0,$$

yielding the coefficients of the orthogonal expansion of

$$\frac{\partial p_n(x,t)}{\partial t} = -\frac{\dot\gamma_n}{\gamma_n}p_n(x,t) - \sum_{k=1}^{m}\alpha_k\dot x_k q_n(x_k,t)\sum_{i=0}^{n}p_{n-i}(x_k,t)p_{n-i}(x,t) \tag{15}$$

$$= -\frac{\dot\gamma_n}{\gamma_n}p_n(x,t) - \sum_{k=1}^{m}\alpha_k\dot x_k q_n(x_k,t)\left[a_n(t)\frac{p_n(x,t)p_{n-1}(x_k,t)-p_n(x_k,t)p_{n-1}(x,t)}{x-x_k} + p_n(x_k,t)p_n(x,t)\right] \tag{16}$$

(Christoffel-Darboux). From the coefficient of p_n:

$$\frac{\dot{\gamma}_n}{\gamma_n} = -\frac{1}{2}\sum_{k=1}^{m} \alpha_k \dot{x}_k q_n(x_k) p_n(x_k)$$

Now, from (14), (2), (3), (4), and (1)

$Wq_n' = (\Omega_n + V)q_n - a_n\Theta_n q_{n-1}, Wp_{n-1}' = a_n\Theta_{n-1}p_n - (\Omega_n + V)p_{n-1}, Wq_{n-1}' = a_n\Theta_{n-1}q_n - (\Omega_n - V)q_{n-1},$

so that, using $p_n q_{n-1} - p_{n-1}q_n = 1/a_n$,

$$\Theta_n(x_k) = 2V(x_k)p_n(x_k)q_n(x_k) = \alpha_k W'(x_k)p_n(x_k)q_n(x_k), \Omega_n(x_k) - V(x_k) = a_n\alpha_k W'(x_k)q_n(x_k)p_{n-1}(x_k), \tag{17}$$

giving (10), (8) follows from $\gamma_{n-1} = a_n\gamma_n$: $\dot{a}_n/a_n = \dot{\gamma}_{n-1}/\gamma_{n-1} - \dot{\gamma}_n/\gamma_n$. Now, (16) becomes

$$\frac{\partial p_n(x,t)}{\partial t} = \frac{\dot{\gamma}_n}{\gamma_n}p_n(x,t) - \sum_{k=1}^{m} \dot{x}_k \, \frac{(\Omega_n(x_k) - V(x_k))p_n(x) - a_n\Theta_n(x_k)p_{n-1}(x)}{W'(x_k)(x - x_k)}. \tag{18}$$

For (9), consider

$$\frac{\partial p_n(x,t)}{\partial t} = \frac{\partial[\gamma_n(t)x^n - \gamma_n(t)(b_0(t) + \cdots + b_{n-1}(t))x^{n-1} + \cdots]}{\partial t} = \frac{\dot{\gamma}_n}{\gamma_n}p_n - \frac{\dot{b}_0 + \cdots + \dot{b}_{n-1}}{a_n}p_{n-1} + \cdots$$

and the coefficient of p_{n-1} in (15):

$$-\frac{\dot{b}_0 + \cdots + b_{n-1}}{a_n} = -\sum_{1}^{m} \alpha_k \dot{x}_k q_n(x_k)p_{n-1}(x_k) = -\frac{1}{a_n}\sum_{1}^{m} \dot{x}_k \frac{\Omega_n(x_k) - V(x_k)}{W'(x_k)} \ .$$

For (11-13), we need the $t-$derivatives of $p_n(x_j(t),t)$, $q_n(x_j(t),t)$, etc.

$dp_n(x_j(t),t)/dt = (\partial p_n(x,t)/\partial t)(x_j) + \dot{x}_j p_n'(x_j)$

$$(3) \Rightarrow \ p_n'(x) = \frac{\Omega_n(x) - V(x)}{W(x)}p_n(x) - a_n\frac{\Theta_n(x)}{W(x)}p_{n-1}(x) = \sum_{k=1}^{m} \frac{(\Omega_n(x_k) - V(x_k))p_n(x) - a_n\Theta_n(x_k)p_{n-1}(x)}{W'(x_k)(x - x_k)},$$

and use (18):

$$dp_n(x_j(t),t)/dt = \frac{\dot{\gamma}_n}{\gamma_n}p_n(x_j,t) + \sum_{\substack{k=1 \\ k\neq j}}^{m} (\dot{x}_j - \dot{x}_k) \ \frac{(\Omega_n(x_k) - V(x_k))p_n(x_j) - a_n\Theta_n(x_k)p_{n-1}(x_j)}{W'(x_k)(x_j - x_k)}, \tag{19}$$

$$\begin{aligned}
\frac{\partial q_n(x,t)}{\partial t} &= \frac{\partial \int_{\mathbb{R}} \dfrac{w(u,t)p_n(u,t)}{x-u}du}{\partial t} \\
&= -\sum_{1}^{m} \alpha_k \dot{x}_k \int_{\mathbb{R}} \frac{w(u)p_n(u)}{(u - x_k)(x - u)}du + \int_{\mathbb{R}} w(u)\frac{\dot{\gamma}_n p_n(u)}{\gamma_n(x-u)}du - \sum_{1}^{m} \dot{x}_k \int_{\mathbb{R}} w(u)\frac{[\Omega_n(x_k) - V(x_k)]p_n(u) - a_n\Theta_n(x_k)p_{n-1}(u)}{W'(x_k)(u - x_k)(x - u)}du \\
&= \frac{\dot{\gamma}_n}{\gamma_n}q_n(x) - \sum_{1}^{m} \dot{x}_k \frac{[\Omega_n(x_k) + V(x_k)](q_n(x) - q_n(x_k)) - a_n\Theta_n(x_k)(q_{n-1}(x) - q_{n-1}(x_k))}{W'(x_k)(x - x_k)}
\end{aligned}$$

$$dq_n(x_j(t),t)/dt = \frac{\dot\gamma_n}{\gamma_n} q_n(x_j,t) + \sum_{\substack{k=1\\k\neq j}}^{m} (\dot x_j - \dot x_k)\, \frac{(\Omega_n(x_k)+V(x_k))q_n(x_j) - a_n\Theta_n(x_k)q_{n-1}(x_j)}{W'(x_k)(x_j-x_k)},$$

and we get (11) as $\alpha_j d(p_n(x_j)q_n(x_j))/dt$ (from (17)). For (12): use (11) with $n-1$ instead of n, use $\Omega_{n-1}(x) = (x - b_{n-1}\Theta_{n-1}(x) - \Omega_n(x)$ from (4), and (12) follows from (10) and also $\sum_1^m \Theta_{n-1}(x_k)/W'(x_k) =$ coefficient of x^{m-1} of $\Theta_{n-1} = 0)$. Finally, we need $dp_{n-1}(x_j(t),t)/dt$ for (13): take (19) with $n-1$, replace $-a_{n-1}p_{n-2}(x_j)$ by $a_n p_n(x_j) - (x_j - b_{n-1})p_{n-1}(x_j)$:

$$dp_{n-1}(x_j(t),t)/dt = -\frac{\dot\gamma_{n-1}}{\gamma_{n-1}} p_{n-1}(x_j,t) - \sum_{\substack{k=1\\k\neq j}}^{m} (\dot x_j - \dot x_k)\, \frac{(\Omega_n(x_k)+V(x_k))p_{n-1}(x_j) - a_n\Theta_{n-1}(x_k)p_n(x_j)}{W'(x_k)(x_j-x_k)}.$$

Exercise: *show that the moment* $\mu_n = \int_{\mathbb{R}} w(u)(u-x_1)^n du$ *of* (6) *enters the solution of the following differential system as* $\mu_n = \nu_{n,1}$,
where $\nu_{n,j} = \int_{\mathbb{R}} w(u)\prod_1^m (u-x_k)^{\beta_k}(u-x_j)^{-1}du$:

$$\dot\nu_{n,j} = \sum_{\substack{k=1\\k\neq j}}^{m} (\dot x_j - \dot x_k)(\alpha_k+\beta_k)\frac{\nu_{n,j}-\nu_{n,k}}{x_j-x_k}, \qquad j=1,\ldots,m$$

with $\beta_1 = n+1, \beta_2 = \ldots = \beta_m = 1$.

This shows that the moments of these generalized Jacobi weights are regular functions of t in any region where the $x_k(t)$'s are distinct. As $a_n^2(t)$ and $b_n(t)$ are ratios of determinants of moments, the movable singular points of their equations can only be poles (the zeros of the denominator determinants).

[1] A.P. MAGNUS, Painlevé-type differential equations for the recurrence coefficients of semi-classical orthogonal polynomials, to appear in *J. Comp. Appl. Math.*

[2] A.P. MAGNUS, Asymptotics for the simplest generalized Jacobi polynomials recurrence coefficients from Freud's equations: numerical explorations. Preprint.